

THE UNITED STATES OF AMERICA

Hioneer Hi-Ared International, Inc.

MACCERS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TILLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR CORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE VE PURPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT ED BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN, FIELD

'PHC77'

In Testimonn Marrest, I have hereunto set my hand and caused the seal of the Plant Anxiety Protection Office to be affixed at the City of Washington, D.C. this twenty-third day of November, in the year two thousand and seven.

Atlest:

Berzu

Commissioner Plant Variety Protection Office Agricultural Marketing Service Syriculture

| REPRODUCE LOCALLY, include form number and | date on all repro | ductions | | | | Form Approved - OMB No. 0581-0055 |
|---|---|--|-------------------------|---|--------------|---|
| U.S. DEPARTME | | | | ne following statements are made in a ne Paperwork Reduction Act (PRA) o | | e with the Privacy Act of 1974 (5 U.S.C. 552a) and |
| SCIENCE AND TECHNOLOGY - P | | | | | | plant variety protection certificate is to be issued |
| APPLICATION FOR PLANT VA (Instructions and information co. | | | , (Ż | U.S.C. 2421). Information is held co | nfidential | until certificate is issued (7 U.S.C. 2426), |
| . NAME OF OWNER | | | 2. | TEMPORARY DESIGNATION OR EXPERIMENTAL NAME | 3. VA | RIETY NAME |
| Pioneer Hi-Bred | l Internatio | nal, Inc. | | | | PHC77 |
| . ADDRESS (Street and No., or R.F.D. No., City, | State, and ZIP Co | ode, and Country) | 5. | TELEPHONE (include area code) | | FOR OFFICIAL USE ONLY |
| 7301 NW | 62 nd Aveni | ıe | | 515/270-4051 | PVPO | NUMBER |
| Johnston | , IA 50131- | 0085 | 6. | FAX (include area code) | - | 200500256 |
| | | | | 515/253-2125 | FILING | S DATE |
| . IF THE OWNER NAMED IS NOT A "PERSON", ORGANIZATION (corporation, partnership, asso | | 8. IF INCORPORATED, GIVE STATE OF INCORPORATION | | DATE OF INCORPORATION | 1 | 11 2005 |
| Corporation | | lowa | . | March 5, 1999 | | May 17, 2005 |
| 0. NAME AND ADDRESS OF OWNER REPRESE | ≓NTATIVE(S) TO | SERVE IN THIS APPLICATION (5) | ret nereni | n listed will receive all paners) | FE | FILING AND EXAMINATION FEES: |
| o, mane and a second of the first field | - | OLIVE IN THIS AT LIGATION. (A) | isi persor | n iisted wili receive ali papers) | E S | \$ 3652.00 |
| _ | | R. Anderson | | | R E | DATE 5/17/05 |
| Rese | | Product Developmen D. Box 85 | t | | E | CERTIFICATION FEE: |
| | | , IA 50131-0085 | | | v E | DATE 10/15/07 |
| · . | , | - | | | D | DATE 10/13/01 |
| TELEPHONE (Include area code) | 12. FAX (Includ | • | ĺ | 13. E-MAIL | on an | derson@pioneer.com |
| 515/270-4051 | | 515/253-2125 | | Stev | GII.AII | derson@pioneer.com |
| 4. CROP KIND (Common Name) Corn | 16, FAMILY N | AME (Botanical) Gramineae | | 18. DOES THE VARIETY CONTA | IN ANY T | RANSGENES? (OPTIONAL) |
| 5. GENUS AND SPECIES NAME OF CROP | 17. IS THE VAI | RIETY A FIRST GENERATION HYB | RID? | ☐ YES ☑ NO IF SO. PLEASE GIVE THE A | SSIGNED | USDA-APHIS REFERENCE NUMBER FOR THE |
| Zea Mays | ☐ YES | ⊠ NO | | APPROVED PETITION TO D COMMERICALIZATION. | EREGUL | ATE THE GENETICALLY MODIFIED PLANT FOR |
| CHECK APPROPRIATE BOX FOR EACH ATTA (Follow instructions on reverse) | ACHMENT SUBM | ITTED | | 20. DOES THE OWNER SPECIFY | | EED OF THIS VARIETY BE SOLD AS A CLASS 3(a) of the Plant Variety Protection Act) |
| a. \(\times \) Exhibit A. Origin and Breeding History of b. \(\times \) Exhibit B. Statement of Distinctness | • | | | ☐ YES (If "yes", answer i | tems 21 a | and 22 below) 🛛 NO (If "no", go to item 23) |
| c. 🗵 Exhibit C. Objective Description of Variety | | | Ī | 21. DOES THE OWNER SPECIFY NUMBER OF CLASSES? | THAT S | EED OF THIS VARIETY BE LIMITED AS TO |
| d. ☐ Exhibit D. Additional Description of the Vee. i. Exhibit E. Statement of the Basis of the Ceeding of t | wner's Ownership | | | ☐ YES ☐ NO | | |
| venification that tissue culture will be depo repository) | | | | | | NDATION REGISTERED CERTIFIED EED OF THIS VARIETY BE LIMITED AS TO |
| g. Filing and Examination Fee (\$3,652), mad States" (Mail to the Plant Variety Protection | le payable to "Tre: on Office) | asurer of the United | | NUMBER OF GENERATIONS | | CED OF THIS VARIETY DE LIVITED AS TO |
| | | | | ☐ YES ☐ NO | | |
| | | | 1 | IF YES, SPECIFY THE NUMB | ER 1,2,3, | etc. FOR EACH CLASS. |
| | | · | | | GISTERE | D CERTIFIED lease use the space indicated on the reverse.) |
| HAS THE VARIETY (INCLUDING ANY HARVES FROM THIS VARIETY BEEN SOLD, DISPOSED | | | | 24. IS THE VARIETY OR ANY CO | MPONEN | IT OF THE VARIETY PROTECTED BY LANT BREEDER'S RIGHT OR PATENT)? |
| OTHER COUNTRIES? ☑ YES □ NO | | | | ☐ YES ⊠ NO | | |
| IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE | | | | IJ YES ⊠ NO IF YES, PLEASE GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED | | |
| IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indicated on reverse.) | | | | IF YES, PLEASE GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED REFERENCE NUMBER. (Please use space indicated on reverse.) | | |
| The owners declare that a viable sample of bas for a tuber propagated variety a tissue culture w | ic seed of the vari /ill be deposited in | ety has been furnished with applicat a public repository and maintained | tion and with for the d | will be replenished upon request in a furation of the certificate. | ccordanc | e with such regulations as may be applicable, or |
| The undersigned owner(s) is(are) the owner of the entitled to protection under the provisions of Sec | his sexually reprod tion 42 of the Pla | duced or tuber propagated plant vari at Variety Protection Act. | iety, and | believe(s) that the variety is new, dis | stinct, unit | form, and stable as required in Section 42, and is |
| Owner(s) is (are) informed that false representat | ion herein can jec | pardize protection and result in pen | alties. | | . • | |
| IGNATURE OF OWNER | | | SIGNAT | URE OF OWNER V | | , , , |
| | | | | Muen | D. | /Inderagon |
| AME (Please print or type) | | | NAME (| Please print or type) | | |
| A CA ANTI-VA CA THE INC. | ···· | | | | | Anderson |
| APACITY OR TITLE | DATE | | | TY OR TITLE | DATE | -16-2005 |
| and the place of the second | | | rxe | esearch Scientist | <u></u> | 10 2000 |
| • | | | | (See reverse for instructions | and inform | afion collection burden statement) |

3ENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner, (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid rariety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 illing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials o make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuanc of the certificates. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvpindex.htm

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and provide evidence that name has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, 10301 Baltimore Avenue, Suite 401 NAL Building, Beltsville, MD 20705. Telephone: (301) 504-5682 http://www.ams.usda.gov/lsg/seed.htm.

TEM

19a.Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;
- (3) evidence of uniformity and stability; and
- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b.Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

United States, Nov. 1, 2004

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's epresentative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any nodification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The alid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing astructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

he U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, plitical beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information stalle, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

o file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD) SDA is an equal opportunity provider and employer.

:T-470 (04-03) designed by the Plant Variety Protection Office using Word 2002.

Exhibit A: Developmental history for PHC77

Pedigree: PH3RC/PH5W4)XA41516X

Pioneer Line PHC77, Zea mays L., a yellow endosperm corn, inbred, was developed by Pioneer Hi-Bred International, Inc. from the single cross hybrid PH3RC (PVP Certificate Number 200200257) X PH5W4 / (PVP Certificate Number 200100244) using the pedigree method of plant breeding. Varieties PH3RC and PH5W4 are proprietary inbred lines of Pioneer Hi-Bred International, Inc. Selfing was practiced from the above hybrid for 7 generations using pedigree selection. During line development, crosses were made to inbred testers for the purpose of estimating the line's combining ability. Yield trials were grown at Princeton, Indiana as well as other Pioneer research locations. After initial testing, additional hybrid combinations have been evaluated and subsequent generations of the line have been grown and hand-pollinated with observations again made for uniformity.

Variety PHC77 has shown uniformity and stability for all traits as described in Exhibit C - "Objective Description of Variety". It has been self-pollinated and ear-rowed 5 generations with careful attention paid to selection criteria and uniformity of plant type to assure genetic homozygousity and phenotypic stability. The line has been increased both by hand and in isolated fields with continued observations for uniformity and stability, and for 3 generations during the final stages of inbred development and seed multiplication. Very high standards for genetic purity have been established morphologically using field observations and using sound lab electrophoresis methodology.

No variant traits have been observed or are expected in PHC77.

The criteria used in the selection of PHC77 were yield, both per se and in hybrid combinations. Late season plant health and late season root lodging, grain quality, stalk lodging resistance, and kernel size were also important criteria considered during selection. Other selection criteria include: Southern Corn Leaf Blight and Giberella Ear Rot tolerance, above average Gray Leaf Spot and Fusarium Ear Rot tolerance.

Exhibit A: Developmental history for PHC77

| Pedigree Grown F0 F0 F1 F2 |
|----------------------------|
| F0 F1 |
| F1 |
| F1 |
| |
| |
| |
| F2 |
| |
| |
| F3 |
| |
| F4 |
| |
| F5 |
| |
| F6 |
| |
| F 7 |
| |
| F8 |
| (Seed) |
| |
| |
| |
| |

^{*}PHC77 was selfed and ear-rowed from F3 through F7 generation.
#Uniformity and stability were established from F5 through F7 generation and beyond when seed supplies were increased.

Exhibit B: Novelty Statement

Variety PHC77 mostly resembles Pioneer Hi-Bred International, Inc. proprietary inbred line PH5W4 (PVP Certificate No. 200100244). Table 1 shows two sample t-tests on data collected primarily in Johnston and Dallas Center, Iowa in 2004. The traits collectively show measurable differences between the two varieties.

Exhibit B: Novelty Statement

Variety PHC77 has a greater plant height (255.3 cm vs 225.1 cm) than variety PH5W4 (Table 1).

Variety PHC77 has a greater ear weight (139.4 g vs 125.6 g) than variety PH5W4 (Table 1).

Variety PHC77 has a light green silk color (Munsell 5GY 9/4) and PH5W4 has a pink silk color (Munsell 10RP 4/8).

Silk color establishes a clear difference between PHC77 and PH5W4.

Exhibit B: Novelty Statement Table(s)

Table 1: Data from Johnston and Dallas Center, Iowa in 2004 presented by trait, across environments, and broken out by environment. Data are supporting evidence for differences between PHC77 and PH5W4. Varieties were grown in 3 locations that had different environmental conditions. Environments had different planting dates and were in different fields. A two-sample t-test was used to compare differences between means.

Plant height (cm)

| Level | Station | Year | Variety-1 | Variety-2 | Cnt-1 | Cnt-2 | Mean-1 | Mean-2 | Mean Diff | StDev-1 | StDev-2 | StErr-1 | StErr-2 | ŭ | t-Wolne | Deck Deel |
|----------------|------------|------|-----------|-----------|-------|-------|--------|--------|-----------|---------|---------|---------|---------|----------|---------|-----------|
| Over All | | | PHC77 | PH5W4 | 15 | 15 | 255.3 | 225.1 | 30.2 | 7.047 | 5.069 | 1.820 | 1 309 | , × | 13.5 | 0000 |
| Station | JHBDA62004 | | PHC77 | PH5W4 | 5 | 'n | 252.4 | 227.2 | 25.2 | 7.797 | 5.404 | 3.487 | 2,417 | 3 ∞ | | 0.000 |
| Station | JHBJH32004 | | PHC77 | PH5W4 | S | Υ | 257.6 | 223.8 | 33.8 | 8.905 | 3.962 | 3.982 | 1.772 | o 00 | 2.5 | 0000 |
| Station | JHYDC22004 | | PHC77 | PH5W4 | 5 | S | 256.0 | 224.4 | 31.6 | 4.000 | 990.9 | 1.789 | 2.713 | · ∞ | 9.7 | 0.000 |
| Ear weight (g) | | | | | | | | | | | | | | | | |
| Level | Station | Year | Variety-1 | Variety-2 | Cnt-1 | Cnt-2 | Mean-1 | Mean-2 | Mean Diff | StDev-1 | StDev-2 | StErr-1 | StErr-2 | T. | t-Value | Prob Pool |
| Over All | | | PHC77 | PH5W4 | 15 | 15 | 139.4 | 125.6 | 13.8 | 9.202 | 13.292 | 2.376 | 3.432 | 28 | 3.3 | 0.003 |
| Environment | JHBDA62004 | | PHC77 | PH5W4 | S | 5 | 140.8 | 122.8 | 18.0 | 4.207 | 17.796 | 1.881 | 7.959 | , ∞ | 2.2 | 0.059 |
| Environment | JHBJH32004 | | PHC77 | PH5W4 | 'n | 5 | 143.6 | 128.0 | 15.6 | 12.973 | 8.367 | 5.802 | 3.742 | | 2.3 | 0.054 |
| Environment | JHYDC22004 | | PHC77 | PH5W4 | ٠, | ν. | 133.8 | 126.0 | 7.8 | 6.834 | 14.646 | 3.056 | 6.550 | ∞ | 1.1 | 0.312 |

Based on previous discussions with the PVP office the traits longitudinal creases and marginal leaf waves were not collected. These traits have low distinguishing power and are variable due to daily fluctuations in water status of the plants. Therefore, we eliminated them from our process based on previous feedback from the PVP office. For insect or disease traits we included data from disease pressure locations only if they were available and paired with the public check. Most often diseases and insect trials are conducted on hybrids since that is the product ultimately sold. In addition, creating consistent disease pressure and infestation levels is costly and difficult.

In cases where less than 15 observations are presented the trait was collected at the plot level as it always has been done in the past. This means many more plants were visually evaluated according to the procedure outlined below, and then a score of the "population" of the plants was recorded for each location.

The experimental design and methods for 2004 were as follows:

The experiment procedures involved three environments with different planting dates per year, planted in 17.42 ft. rows with 2 rows for each variety. Approximately 24-30 plants emerged in each of 2 rows for a total of around 48 to 60 plants being evaluated at each location and 144 to 180 plants across locations. For plant level traits, we sampled 5 representative plants from the 2 rows of the 2 row plot (group) of plants at each location. For plot level traits we evaluated the 2 row plot (group) and gave a representative score or average on the 48-60 plants in the group within an experiment.

Some traits can be especially variable under different environmental factors influenced by weather, soil type, or planting dates. Varying temperatures or day length could impact the meristem growth during various tissue differentiation stages. The meristem differentiation of the ear and other tissues could be impacted as well as the success of pollination during flowering and frequency of kernel abortion during grain fill.

We have included weather data in the table that follows.

| | SKOW S | ING DEGR | GROWING DEGREE UNITS (GDU'S) | GDU's) | ā | PRECIPITATION (inches) | 10N (inche | 1 |
|-----------|-----------|----------|------------------------------|----------|---------|------------------------|------------|---------------------------------------|
| | 20 | 2003 | 20 | 2004 | 20 | 2003 | 100 | 2004 |
| | D. Center | Johnston | D. Center | Johnston | D. Cent | Johnston | Cent | Pohneton |
| May | 375 | 380 | 548 | 527 | 5.7 | 5 43 | T | 31" |
| June | 909 | 604 | 609 | 610 | 1.92 | 4 23 | 1 97 | 200 |
| July | 628 | 782 | 723 | 736 | 0.18 | 3.4 | 2 20 | 200 |
| August | 795 | 786 | 612 | 615 | 0.44 | 0.51 | 1 95 | 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 |
| September | 456 | 468 | 598 | 560 | 2.19 | 2.52 | 38 | 5 6 |
| TOTAL | 2860 | 3020 | 3090 | 3048 | 10.43 | 16.09 | 14.78 | 24 87 |

Calculate GDU's
Growing Degree Units use following formula: GDU = ((T1+T2)/2)-50
Where T1 = minimum temperature for a given day with 80 degrees Fahrenheit as the minimum temperature used and 86 degrees Fahrenheit is the maximum temperature used.
Where T2 = maximum temperature for a given day with 86 degrees Fahrenheit as the maximum temperature used and 50 degrees Fahrenheit is the minimum temperature used.

United States Department of Agriculture, Agricultural Marketing Service Science and Technology, Plant Variety Protection Office National Agricultural Library Building, Room 400 Beltsville, MD 20705-2351

OBJECTIVE DESCRIPTION OF VARIETY CORN (Zea mays L.)

| Pioneer Hi-Bred Internat | ional, Inc | l Variety Seed S | Source | I Variety Name or 1 I PHC77 | Femporary Designation |
|--|---|---|---|---|---|
| Address (Street & No., or 7301 NW 62nd Avenue, I | R.F.D. No., City, State, Zip P.O. Box 85, Johnston, Io | Code and Country wa 50131-0085 | FOR OFFICIAL | | PVPO Number |
| adding leading zeroes if n | nber that describes the vari ecessary. Completeness si an adequate variety descri | hould be striven for to e | establish an adequate vari | e spaces below. Right | justify whole numbers by designated by a "*" are |
| 01. Light Green 02. Medium Green 03. Dark Green 04. Very Dark Green 05. Green-Yellow | n conjunction with Munsell 06. Pale Yellow 07. Yellow 08. Yellow-Orange 09. Salmon 10. Pink-Orange | 11. Pink 12. Light Red 13. Cherry Red 14. Red 15. Red & White | 16. Pale Purple17. Purple18. Colorless19. White20. White Capped | 21. Buff 22. Tan 23. Brown 24. Bronze 25. Variegated (Des | 26. Other (Describe) |
| Family Meml B14 CM10 B37 B37 B73 N192 C103 M017 Oh43 A619 | | ar (in background and n ellow Dent (Unrelated): Co109, ND246 Oh7, T232 W117, W153R W182BN White Dent: Cl66, H105, Ky22 | | Sweet Corn: C13, Iowa512 Popcorn: | 25, P39, 2132 722, HP301, HP7211 |
| 1. TYPE: (describe interm 3 (1=Sweet, 2=D Flint-Dent | nediate types in "Comment Dent, 3=Flint, 4=Flour, 5=P | s" section) op, 6=Ornamental, 7=F | Pipecorn) | I Standard Inbred I I 2 Type | Name B73 |
| 2. REGION WHERE DEV 3 (1=N.West, 2= | ELOPED IN THE U.S.A.: N.Central, 3=N.East, 4=S. | East, 5=S.Central, 6=S | .West, 7=Other | I Standard Seed S | ource PI 550473 |
| DAYS HEAT 68 1,4 | 465.7 From emergence to 145.5 From emergence to 46 From 10% to 90% p From 50% silk to op | 50% of plants in silk 50% of plants in pollen ollen shed | · | DAYS 65 64 | HEAT UNITS 1,382.5 1,366.8 49 |
| 14.5 cm Length of 0.0 Average Numl 1.1 Average Numl | to base of top ear node) Top Ear Internode | <u>1</u> | Dev. Sample Size 1.89 30 9.45 30 2.19 30 0.03 6 0.20 6 =Dark | I <u>238.2</u> I <u>98.1</u> I <u>15.5</u> I <u>0.0</u> | St.Dev. Sample Size 19.92 30 16.92 30 1.68 30 0.02 6 0.07 6 |
| Application Variety Data | | P | age 1 | I Standard Inbred I | Data |

| Application Variety Data 5. LEAF 9.1 cm Width of Ear Node Leaf 85.7 cm Length of Ear Node Leaf 6.8 Number of leaves above top ear 21.8 Degrees Leaf Angle (Measure from 2nd leaf above ear at anthesis to stalk above 4 Leaf Color (Munsell Code) 7.5GY34 3 Leaf Sheath Pubescence (Rate on scale from 1=none to 9= Marginal Waves (Rate on scale from 1=none to 9=many) Longitudinal Creases (Rate on scale from 1=none to 9=man) 5. TASSEL: 4.4 Number of Primary Lateral Branches 4.3.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy 6 Anther Color (Munsell Code) 7.5Y86; | St.Dev. 0.84 5.22 0.89 6.44 leaf) like peach from the second se | Sample Size 30 30 30 30 30 30 30 3 | Mean | St.Dev. 0.79 3.32 0.77 5.08 Code) 7.50 St.Dev. 1.30 8.41 | Sample Si |
|--|--|--|--|--|---------------------------------|
| 9.1 cm Width of Ear Node Leaf 85.7 cm Length of Ear Node Leaf 6.8 Number of leaves above top ear 21.8 Degrees Leaf Angle (Measure from 2nd leaf above ear at anthesis to stalk above 4 Leaf Color (Munsell Code) 7.5GY34 3 Leaf Sheath Pubescence (Rate on scale from 1=none to 9= Marginal Waves (Rate on scale from 1=none to 9=many) Longitudinal Creases (Rate on scale from 1=none to 9=man) 5. TASSEL: 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | 0.84 5.22 0.89 6.44 leaf) slike peach from the second | 30 30 30 30 30 30 30 30 | 8.7 80.9 6.4 22.4 4 (Munsell (3 — — — — Mean 7.3 15.6 | 0.79 3.32 0.77 5.08 Code) 7.50 St.Dev. 1.30 | SY3 <mark>4</mark> Sample Si |
| 85.7 cm Length of Ear Node Leaf 6.8 Number of leaves above top ear 21.8 Degrees Leaf Angle (Measure from 2nd leaf above ear at anthesis to stalk above 4 Leaf Color (Munsell Code) 7.5GY34 3 Leaf Sheath Pubescence (Rate on scale from 1=none to 9= Marginal Waves (Rate on scale from 1=none to 9=many) Longitudinal Creases (Rate on scale from 1=none to 9=man) 5. TASSEL: 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | 0.84 5.22 0.89 6.44 leaf) slike peach from the second | 30 30 30 30 30 30 30 30 | 8.7 80.9 6.4 22.4 4 (Munsell (3 — — — — Mean 7.3 15.6 | 0.79 3.32 0.77 5.08 Code) 7.50 St.Dev. 1.30 | SY3 <mark>4</mark> Sample Si |
| 6.8 Number of leaves above top ear 21.8 Degrees Leaf Angle (Measure from 2nd leaf above ear at anthesis to stalk above 4 Leaf Color (Munsell Code) 7.5GY34 3 Leaf Sheath Pubescence (Rate on scale from 1=none to 9= Marginal Waves (Rate on scale from 1=none to 9=many) Longitudinal Creases (Rate on scale from 1=none to 9=man) 5. TASSEL: 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy) | 0.89 6.44 leaf) like peach f y) St.Dev. 1.25 20.52 3.68 | 30 30 30 30 30 30 30 30 | 80.9 6.4 22.4 4 (Munsell (3 — — — — Mean 7.3 15.6 | 3.32 0.77 5.08 Code) 7.50 St.Dev. 1.30 | SY3 <mark>4</mark> Sample Si |
| 21.8 Degrees Leaf Angle (Measure from 2nd leaf above ear at anthesis to stalk above 4 Leaf Color (Munsell Code) 7.5GY34 3 Leaf Sheath Pubescence (Rate on scale from 1=none to 9= Marginal Waves (Rate on scale from 1=none to 9=many) Longitudinal Creases (Rate on scale from 1=none to 9=man) 5. TASSEL: 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | 0.89 6.44 leaf) like peach f y) St.Dev. 1.25 20.52 3.68 | 30 30 1 1 1 1 1 1 1 1 1 | 6.4 22.4 4 (Munsell 0 3 — — — Mean 7.3 15.6 | 0.77 5.08 Code) 7.50 St.Dev. 1.30 | SY3 <mark>4</mark> Sample Si |
| (Measure from 2nd leaf above ear at anthesis to stalk above 4 Leaf Color (Munsell Code) 7.5GY34 3 Leaf Sheath Pubescence (Rate on scale from 1=none to 9= Marginal Waves (Rate on scale from 1=none to 9=many) Longitudinal Creases (Rate on scale from 1=none to 9=man 5. TASSEL: 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | leaf) like peach filling) St.Dev. 1.25 20.52 3.68 | 30 | 4 (Munsell (3 ——————————————————————————————————— | 5.08 Code) 7.50 St.Dev. 1.30 | SY3 <mark>4</mark> Sample Si |
| 4 Leaf Color (Munsell Code) 7.5GY34 3 Leaf Sheath Pubescence (Rate on scale from 1=none to 9= | St.Dev. 1.25 20.52 3.68 | Sample Size 1 | 4 (Munsell (3 ———————————————————————————————————— | St.Dev. | SY3 <mark>4</mark> Sample Si |
| 4 Leaf Color (Munsell Code) 7.5GY34 3 Leaf Sheath Pubescence (Rate on scale from 1=none to 9= | St.Dev. 1.25 20.52 3.68 | Sample Size 1 30 1 30 i | 3 Mean 7.3 15.6 | St.Dev. 1.30 | Sample Si |
| Marginal Waves (Rate on scale from 1=none to 9=many) Longitudinal Creases (Rate on scale from 1=none to 9=man TASSEL: 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | St.Dev. 1.25 20.52 3.68 | Sample Size 1 30 1 30 i | 3 Mean 7.3 15.6 | St.Dev. 1.30 | Sample Si |
| Marginal Waves (Rate on scale from 1=none to 9=many) Longitudinal Creases (Rate on scale from 1=none to 9=man 5. TASSEL: | St.Dev. 1.25 20.52 3.68 | Sample Size 1 30 1 30 i | — — Mean <u>7.3</u> 1 <u>5.6</u> | <u>1.30</u> | |
| 6. TASSEL: 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | St.Dev. 1.25 20.52 3.68 | 30 I 30 I | <u>7.3</u> <u>15.6</u> | <u>1.30</u> | |
| 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | 1.25 20.52 3.68 | 30 I 30 I | <u>7.3</u> <u>15.6</u> | <u>1.30</u> | |
| 4.4 Number of Primary Lateral Branches 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | 1.25 20.52 3.68 | 30 I 30 I | <u>7.3</u> <u>15.6</u> | <u>1.30</u> | |
| 43.7 Degrees Branch Angle from Central Spike 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | 20.52 3.68 | <u>30</u> 1 | <u>15.6</u> | | 2 |
| 62.8 cm tassel Length (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | 3.68 | | | 0.41 | |
| (from top leaf collar to tassel tip) 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | | <u>30</u> 1 | 55.4 | | <u>.</u> <u>.</u> |
| 4 Pollen Shed (Rate on scale from 0=male sterile to 9=heavy | shed) | | | <u>4.16</u> | 2 |
| | sileu) | 1 | E | | |
| | |] | <u>5</u> 7 (Munsell 0 | Code) <u>5Y8.</u> | ε l |
| 12 Glume Color (Munsell Code) 7.5RP48 | | : | 7 (Munseil (| | |
| 1 Bar Glumes (Glume Bands): 1=Absent, 2=Present | | i | <u>2</u> (Mansen (| Joue) <u>331</u> | 30 |
| a. EAR (Unhusked Data): | | · . I | | | |
| 1 Silk Color (3 days after emergence) (Munsell Code) | 5GY | /9Å | 1 Munsell C | ode 2.5G | Y94 |
| 2 Fresh Husk Color (25 days after 50% silking) (Munsell Code | e) <u>5GY</u> | <u>768</u> , 1 | | ode 5GY | |
| 21 Dry Husk Color (65 days after 50% silking) (Munsell Code) | 2.51 | 78.54 | 21 Munsell C | | |
| 1 Position of Ear at Dry Husk Stage: 1=Upright, 2=Horizontal, | , 3=Pendent | , | | | - i |
| 7 Husk Tightness (Rate on scale from 1=very loose to 9=very | tight | ľ | 8 | | |
| 2 Husk Extension (at harvest): 1=Short(ears exposed), 2=Med | dium (<8cm) |), 3=Long | 3 8 3 | | |
| (8-10cm beyond ear tip), 4=Very Long (>10cm) | | - 1 | _ | | |
| b. EAR (Husked Ear Data) | St. Dev. | Sample Size I | Mean | St.Dev. | Sample Siz |
| <u>14.6</u> cm Ear Length | 0.94 | 30 1 | 13.1 | 1.09 | ` a |
| 42.7 mm Ear Diameter at mid-point | 1.89 | <u>30</u> 1 | 44.6 | 3.23 | 3 |
| 130.3 gm Ear Weight | 18.53 | <u>30</u> l | 118.8 | 43.50 | 102 |
| 14.8 Number of Kernel Rows | 1.24 | 30 I | <u>17.9</u> | 1.74 | 3 |
| 2 Kernel Rows: 1=Indistinct, 2=Distinct | <u></u> | <u> </u> | 2 | 1111 | 2 |
| 2 Row Alignment: 1=Straight, 2=Slightly Curved, 3=Spiral | | i | 2 | | |
| 7.6 cm Shank Length | 0.93 | <u>30</u> i | 7 1 | <u>2.11</u> | 3 |
| 2 Ear Taper: 1=Slight cyl., 2=Average slightly con., 3=Extreme | e conical | i | 2 2 7.4 2 | 4.11 | _ |
| KERNEL (Dried): | St.Dev. | Sample Size I | Mean | St.Dev. | Sample Siz |
| 11.0 mm Kernel Length | 0.76 | 30 | 11.0 | 1.61 | Sumple Ca |
| 7.7 mm Kernel Width | 0.60 | <u>30</u> i | 6.5 | 0.68 | ŝ |
| 4.5 mm Kernel Thickness | 0.68 | <u>30</u> I | <u>4.1</u> | 0.48 | 5 |
| 26.9 % Round Kernels (Shape Grade) | 5.62 | <u>50</u> i | 10.1 | 5.29 | 3 |
| 1 Aleurone Color Pattern: 1=Homozygous, 2=Segregating (de | scribe) | <u>⊻</u> ' | 1 (describe) | | |
| 7 Aleurone Color (Munsell Code) 10YR612 | oonbe, | i | 7 Munsell C | | 5Y812 |
| 7 Hard Endosperm Color (Munsell Code) 10YR610 | | i | 7 Munsell C | | YR7/12 |
| 3 Endosperm Type: 1=Sweet(su1), 2=Extra Sweet(sh2), 3=No | rmal Starch | 4=High | 3 (describe) | | 1111112 |
| Amylose Starch, 5=Waxy Starch, 6=High Protein, 7=High Lys | sine 8=Suna | r Sweet | ∑ (describe) | | |
| (se), 9=High Oil, 10=Other | ο, υ-οαρε | . Sweet 1 | | | |
| 28.0 gm Weight per 100 kernels (unsized sample) | 2.97 | 1 <u>6</u> 1 | 20.3 | <u>7.63</u> | |
| COB: | St Day | Comple Circ. | NA+ | C4 D | Cana-l- Ci |
| 23.4 mm Cob Diameter at mid-point | St.Dev. <u>1.52</u> | Sample Size I 30 I | Меап 27.5 | St.Dev 1.41 | Sample Si |
| 14 Cob Color (Munsell Code) 2,5YR3B | 1,04 | <u>55</u> 1 | <u>11.9</u> 11 Munsell | | 0R66 |
| <u> </u> | | | 11 Manaell | | <u> </u> |
| pplication Variety Data | Page 2 | I | Standard Inbred Da | ata | |

Note: Use chart on first page to choose color codes for color traits.

| 11 INSECT RESISTANCE | CE (Rate from 1(most susceptible) to | 0 (most resistant): Legye blank | |
|---|---|--|--|
| if not tested | DE (Rate from Timost susceptible) ti | o 9 (most resistant); Leave blank St. Dev. Sample Size | |
| | ite (Oligonychus pratensis) | ot. Dev. Sample Size | |
| Corn Earworm (Helic | everna zee) | | l Banks Grass Mite |
| Leaf Feeding | overpa zea) | | Corn Earworm |
| | man lamed set | • | Leaf Feeding |
| Sik reeding _ | mg larval wt. | | · |
| _ Ear Damage | * / **** | | L Ear Damage |
| | f (Rhopalosiphum maidis) | | Corn Leaf Aphid |
| | e (Carpophilus dimidiatus) | | l Corn Sap Beetle |
| European Corn Bore | · (Ostrinia nubilalis) | | I European Corn Borer |
| | Typically Whorl Leaf Feeding) | | I 1 st Generation |
| 2 nd Generario | r (Typically Leaf Sheath-Collar Feed | ing) | I _ 2 nd Generation |
| Stalk Tunneling: | cm tunneled/plant | | · . |
| Fall Armyworm (Spoo | loptera frugiperda) | | l Fall Armyworm |
| _ Leaf-Feeding | , | | Leaf-Feeding |
| | mg larval wt. | | |
| Maize Weevil (8 | | | Maize Weevil |
| | ntopinius zeamais, | | _ Mai28 VIGGVII |
| Northern Poots | orm (Diabrotica barberi) | | I Northorn Poetworm |
| | | | Northern Rootworm |
| Southern Rootw | rorm (Diabrotica undecimpunctata) | | Southern Rootworm |
| | orer (Diatraea grandiosella) | | I Southwestern Corn Borer |
| _ Leaf Feeding | | | Leaf Feeding |
| Stalk Tunneling: | cm tunneled/plant | | <u> </u> |
| | ider Mite (Tetranychus urticae) | | I _ Two-spotted Spider Mite |
| _ Western Rootwe | orm (Diabrotica virgifera virgifera) | | I _ Western Rootworm |
| _ Other (Specify)_ | | | l Other (Specify) |
| % Dropped Ears _ % Pre-anthesis % Pre-anthesis % Post-anthesis | 55 days after anthesis) (Rate on sca (at 65 days after anthesis) Brittle Snapping Root Lodging : Root Lodging (at 65 days after anti | nesis) | 2 Stay Green No Dropped ears No Pre-anthesis Brittle Snapping No Pre-anthesis Root Lodging No Pre-anthesis Root Lodging No Pre-anthesis Root Lodging |
| 5,572.0 Kg/na fi | eld of Inbred Per Se (at 12-13% grai | n moisture) | 5.714.0 Yield |
| 13. MOLECULAR MARK | ERS: (0=data unavailable; 1=data av | /ailable but not supplied; 2=data | supplied.) |
| 1 Isozymes | _ RFLP's | _ RAPD's | Other (Specify) |
| REFERENCES: | | | |
| | 6 (1 6) 16 (1 4 4 4 1 1 | | |
| Emerson, R.A., G.W. Bea | em for the Classification of Corn Inb Idle, and A.C. Fraser, 1935. A sumn Chamuris, A.Y. Rossman, 1989. F | nary of Linkage Studies in Maize. | |
| • | | | · |
| Jugenheimer, R.W. 1976. McGee, D.C. 1988. Maize Munsell Color Chart for Pi The Mutants of Maize. 19 Shurtleff, M.C. 1980. Com | orn: Culture, Processing, Products. A Corn: Improvement, Seed Production Diseases. APS Press, St. Paul, MN ant Tissues. Macbeth. P.O. Box 230 68. Crop Science Society of America Opendium of Corn Diseases. APS Production (Editors). 1988. Corn and Co | on, and Uses. John Wiley & Son: J. 150 pp. J. Newburgh, N.Y. 12551-0230 a. Madison, WI. ess, St. Paul, MN. 105 pp. | |
| | ored Lines of Ohio A.E.S., Bul. 831. ulture 1936, 1937. Yearbook. | 1959. | |

COMMENTS (e.g. state how heat units were calculated, standard inbred seed source, and/or where data was collected. Continue in Exhibit D)

Insect, disease, brittle snapping and root lodging data are collected mainly from environment where variability for the trait can be obtained within the experiment.

CLARIFICATION OF DATA IN EXHIBITS B AND C

Please note the data presented in Exhibit B and C, "Objective Description of Variety," are collected primarily at Johnston and Dallas Center, Iowa. The data in Table 1 are from two sample t-tests using data collected in Johnston and Dallas Center, IA. These traits in exhibit B collectively show distinct differences between the two varieties.

| REPRODUCE | LOCALLY. Include form number and edition date on all reproduction | | -04 00FF | | |
|--|--|---|---|--|--|
| | U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E EMENT OF THE BASIS OF OWNERSHIP | Application is required in order to determ certificate is to be issued (7 U.S.C. 2421) confidential until the certificate is issued (| ine if a plant variety protection | | |
| | ER HI-BRED INTERNATIONAL, INC. | 2.TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER | 3. VARIETY NAME PHC77 | | |
| 4 .ADDRESS | (Street and No., or R.F.D. No., City, State, and ZIP, and Country) | 5. TELEPHONE (include area code) | 6. FAX (include area code) | | |
| 7301 N | W 62 nd AVENUE | 515-270-4051 | 515-253-2125 | | |
| P.O.BC | | 7.PVPO NUMBER | | | |
| JOHNS | TON, IA 50131-0085 | 200500 | | | |
| 8.Does the app | olicant own all rights to the variety? Mark an "X" in the appropriate | e block. If no, please explain: 🛛 YES | □NO | | |
| | | ર્ય લ- | | | |
| | | ~ ? | | | |
| 9.Is the applica | ant (individual or company) a U.S. national or a U.S. based compa | any? If no, give name of country, X | ES F1 NO | | |
| | , a constant of a constant con | any. If no, give name of country. | | | |
| 10. Is the appli | cant the original owner? | answer <u>one</u> of the following: | | | |
| a. If the | original rights to variety were owned by individual(s), is (are) the | original owner(s) a U.S. National(s)? | | | |
| • | ☐ YES ☐ NO If no, give name of country | | | | |
| b. If the | original rights to variety were owned by a company(ies), is (are) | the original owner(s) a U.S. based company | ? | | |
| | | | | | |
| 11. Additional e | explanation on ownership (Trace ownership from original breeder | to current owner. Use the reverse for extra s | space if needed): | | |
| is the emploration | fi-Bred International, Inc. (PHI), Des Moines, Iowa, and/or its who ployer of the plant breeders involved in the selection and develop on has the sole rights and ownership of PHC77 pursuant to writte ety was created. No rights to this variety are retained by any indiv | ment of PHC77. Pioneer Hi-Bred Internation contracts that assign all rights in the variet | nal and/or Pioneer Overseas | | |
| PLEASE NOTE | : | | | | |
| Plant variety prot | ection can only be afforded to the owners (not licensees) who meet the | following criteria: | | | |
| If the rights which affore | to the variety are owned by the original breeder, that person must be a ds similar protection to nationals of the U.S. for the same genus and sp | a U.S. national, national of a UPOV member coecies. | ountry, or national of a country | | |
| 2. If the rights member con | to the variety are owned by the company which employed the original untry, or owned by nationals of a country which affords similar protect | breeder(s), the company must be U.S. based, of tion to nationals of the U.S. for the same genus | owned by nationals of a UPOV and species. | | |
| 3. If the applic | ant is an owner who is not the original owner, both the original owner | and the applicant must meet one of the above co | riteria. | | |

The original breeder/owner may be the individual or company who directed the final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definitions.

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 0.1 hour per response, including the time for reviewing the instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal employment opportunity provide and employer.